

## Alcohol Consumption

**Table 9. Percent (standard error) of women who reported drinking alcohol during the 3 months before pregnancy. Ques. # 32a**

Maternal Characteristics	percent (s.e.)
<b>Total</b>	41.4(1.3)
<b>Race</b>	
Black	22.9(1.8)
White	54.2(1.7)
<b>Age</b>	
Less than 20	23.5(2.9)
20-29	42.4(1.7)
30+	48.7(2.5)
<b>Education</b>	
Less than High School	27.9(2.7)
High School	38.3(2.1)
More than High School	51.0(2.0)
<b>Marital Status</b>	
Married	47.7(1.7)
Other	33.4(1.9)
<b>Birthweight</b>	
LBW (<2500 grams)	32.2(3.5)
NBW (>= 2500 grams)	42.1(1.4)
<b>Medicaid Status*</b>	
Medicaid	34.3(1.7)
Other	49.9(1.9)

**Table 10. Percent (standard error) of women who reported drinking alcohol during the last trimester of pregnancy, according to sociodemographic characteristics. Ques. # 33a**

Maternal Characteristics	percent (s.e.)
<b>Total</b>	5.2(0.6)
<b>Race</b>	
Black	3.8(0.9)
White	6.1(0.8)
<b>Age</b>	
Less than 20	2.8(1.2)
20-29	4.0(0.7)
30+	8.6(1.5)
<b>Education</b>	
Less than High School	4.6(1.3)
High School	4.4(0.9)
More than High School	5.9(1.0)
<b>Marital Status</b>	
Married	4.9(0.8)
Other	5.1(0.9)
<b>Birthweight</b>	
LBW (<2500 grams)	5.5(1.6)
NBW (>= 2500 grams)	5.0(0.6)
<b>Medicaid Status*</b>	
Medicaid	4.9(0.8)
Other	5.2(0.9)

\* A mother is considered "on Medicaid" if she was enrolled in Medicaid during any of the following time periods: just before pregnancy, during prenatal care or during delivery.